

**Canine Lymphosarcoma-
Short Madison Wisconsin Protocol**

Patient Label

Weight at start of therapy _____

BSA _____

Week	Date	Weight	Drug	Dose	Amount given / location
1			Vincristine	0.5 mg/m ² IV	
			Prednisone	2 mg/kg PO daily	
2			Cyclophosphamide	250 mg/m ² IV or PO	
			Prednisone	1.5 mg/kg daily	
3			Vincristine	0.7 mg/m ² IV	
			Prednisone	1.0 mg/kg daily	
4			Doxorubicin	30 mg/m ² IV	
			Prednisone	0.5 mg/kg	
6			Vincristine	0.7 mg/m ² IV	
7			Cyclophosphamide	250 mg/m ² IV or PO	
			Prednisone	0.5 mg/kg day of tx	
8			Vincristine	0.7 mg/m ² IV	
9			Doxorubicin	30 mg/m ² IV	
11			Vincristine	0.7 mg/m ² IV	
13			Cyclophosphamide	250 mg/m ² IV or PO	
			Prednisone	0.5 mg/kg day of tx	
15			Vincristine	0.7 mg/m ² IV	
17			Lomustine	60 mg/m ² PO	
19			Vincristine	0.7 mg/m ² IV	
21			Cyclophosphamide	250 mg/m ² IV or PO	
			Prednisone	0.5 mg/kg day of tx	
23			Vincristine	0.7 mg/m ² IV	
25			Doxorubicin	30 mg/m ² IV	

Always check CBC before each chemotherapy treatment. Neutrophils should be > 2000 or chemotherapy should be delayed 1 week when possible.

All IV chemotherapy drugs must go IV!!!

Owners must use caution when administering oral chemotherapeutics.